

Nickel and cadmium concentrations in the coral family, Faviidae and surrounding sediments in the south of Qeshm Island, The Persian Gulf

Alireza Riahi Bakhtiari^{1*}

Zahra Darvishnia²

Ehsan Kamrani³

Mir Masoud Sajjadi⁴

Maryam Fakharzadeh⁵

Zahra Biabannavard⁶

1. Assistant Professor of Department of Environmental Science, of Natural Resources and Marine Sciences Faculty, Tarbiat Modares University, Noor, Mazandara, Iran.

2. 6. MSC graduated of Marine Biology, University of Hormozgan, Bandar Abbas, Iran

3. Associate Professor of Department of Marine Biology, Faculty of Science, University of Hormozgan, Bandar Abbas, Iran

4. Associate Professor of Fisheries Department, Faculty of Natural Resources, University of Guilan, Some-Sara, Iran

5. M.Sc. Student of Aquaculture, Persian Gulf University, Bushehr, Iran

***Corresponding author:**

ariyahi@gmail.com

Received date: 2015/08/04

Reception date: 2015/09/02

Abstract

The following research was conducted to determining the nickel (Ni) and cadmium (Cd) concentrations in skeleton of coral family (Faviidae) and sediments of dominant in Zeyton Park, Naz Islands and Shib-deraz stations, south of Qeshm Island -Persian Gulf in summer 2012. The analysis of these heavy metal levels in digested samples of skeleton and sediments was done using graphite furnace Atomic Absorbtion Spectrophotometry (SHIMADZU, AA 670G) for Cd and Flame Atomic Absorption Spectrophotometry (SHIMADZU, AA 670) Ni. The results indicated that there were significant differences in concentrations of Ni and Cd in skeleton of coral family (Faviidae) among Zeyton Park, Naz Islands Shib-deraz stations ($P<0.001$) and concentration of Ni and Cd in sediments of coral family (Faviidae) there are significant differences between Zeyton Park, Naz Islands and Shib-deraz stations ($P<0.05$ and $P<0.001$, respectively). There was a negative correlation between Ni concentration in Faviidae coral and its surrounding sediments in Shib-deraz stations. Besides, there was statistical correlation in Cd concentration between coral family (Faviidae) in Zeyton Park and Shib-deraz stations. Coral Skeletons and sediments in the study area was not contaminated with heavy metals Ni and Cd compared to the results with universal standards. So, concerning the immense importance of coral ecosystem in the southern Qeshm Islands and the result of this research, corals can be suggested as a pollution biomonitor.

Keywords: Bioaccumulation, Heavy metals, Coral, Qeshm Islands, Persian Gulf.