## The crude oil biodegradation by *Fisherella* (*Fischerella* sp. ISC67) and *Nostoc* (*Nostoc* sp. ISC101) Cyanobacteria

Leila Nemati<sup>1</sup> Arezoo Tahmourespour<sup>2\*</sup> Neda Soltani<sup>3</sup>

1. MSc. Soil Sciences Department, Isfahan (Khorasgan) Branch, Islamic Azad University, Isfahan, Iran

2. Associate Prof, Basic Medical Sciences Department, Isfahan (Khorasgan) Branch, Islamic Azad University, Isfahan, Iran 3. Associate Prof. Department of Biology, ACECR, Research Institute of Applied Science, University Shahid Beheshti, Tehran. Iran

\*Corresponding author: atahmoures@khuisf.ac.ir

Received date: 2019.10.08 Reception date: 2020.0116

## Abstract

The purpose of this study was to investigate the growth and potential of crude oil biodegradation by Fischerella and Nostoc. In this study (2015), the growth rate of cyanobacterial species such as Fischerella sp. ISC107 and Nostoc sp. ISC101 were determined using chlorophylla content measurement at 663nm under different treatments of crude oil (control, 0.1, 0.2, 0.4, 0.8 and 1.6%). The crude oil concentration was analyzed by gas chromatography (GC-FID) after 21 days, and also the biodegradation rate was determined. Results showed that by increasing the concentration of crude oil, the amount of chlorophyll-a was significantly decreased. So, the growth rate (based on chlorophyll-a) of these cyanobacteria were almost nearly equal or lower than the control sample in the presence of crude oil. The maximum and minimum percentages of crude oil biodegradation, in the treatments of 0.1 and 1.6% were 88.27% and 44.72% (Fischerella) and 91% and 47.46% (Nostoc), respectively after 21 days. The rate of biodegradation in all treatments was significantly (P<0.05) higher than control. According to results of this study, Fischerella ISC107 and Nostoc ISC 101 were shown great potential for crude oil biodegradation and can be used as valuable strains for biodegrading, elimination or the reduction of oil pollution in contaminated areas.

**Keywords:** Crude oil; Biodegradation; *Fischerella*; *Nostoc*; Chlorophyll-a.