Seasonal Diversity and Frequency of Planktonic Bacillariophyta (Diatoms) in Coastal Waters of the Southern Caspian Sea

Tayebeh Omidmoazam¹ Maryam Akhoundian^{2, 3*} Maryam Fallahi Kapoorchali⁴ Shila Omidzahir^{2, 3}

1. MSc, Department of Marine biology, Faculty of Marine science, University of Mazandaran, Babolsar, Iran 2. 4. Assistant professor, Department of Marine biology, Faculty of Marine science, University Mazandaran, of Babolsar, Iran 3. Professor, Inland Water Center, Aquaculture Iranian Fisheries Science Research Agriculture Institue (IFSRI), Research Education and extension

Organization (AREEO). Bandar-e Anzali. Iran

*Corresponding author:

m.akhoundian@umz.ac.ir

Received date: 2020.01.16 Reception date: 2020.03.05

Abstract

This study aimed to investigate the biodiversity and abundance of diatoms in coastal waters of Sari and Babolsar on 2015, in two linear transects and 8 stations from estuary to 20 m depth. Sampling the water and measuring some environmental parameters such as temperature, salinity, acidity and dissolved oxygen, followed by evaluation of diatoms (Bacillariophyta) qualitatively and quantitatively. Margalef, Pielo and Shannon-wiener indices were used to assess richness, evenness and diversity of diatoms, respectively. There was no correlation between environmental parameters and diversity and abundance of diatoms, but only salinity; also, the mean of diatom diversity indices in both season was higher in Babolrood estuary (1.20 and 0.79). Also, the mean diversity indics of diatoms was higher in winter (0.62) than in summer (0.41). In this study, 17 genera of diatoms were identified and counted; and according the results, Cyclotella sp. was the most abundant diatom in both seasons, and Surrirella sp. was found only in estuaries. Ecological indices' assay showed that, this area is suffering of a kind of ecological disturbance, which indicated that the risk of ecological disturbance in the study area, is more serious than previous research.

Keywords: Diversity, Abundance, Diatoms, Bacillariophyta, ecological indices, Caspian Sea.