Chronic and acute effects of nickel and cobalt on chlorophyll a and carotenoids of marine micro algae *Nannochloropsis oculata*

Morteza Beygi Malekabadi¹ Rasool Zamani-Ahmadmahmoodi*² Ruhollah Rahimi² Hajar Ebrahimi³

- 1. Master Science Student in Fisheries Science, Faculty of Natural Resources and Earth Science, Shahrekord University, Shahrekord, Iran
- 2. Department of Fisheries and Environmental Sciences, Faculty of Natural Resources and Earth Science, Shahrekord University, Shahrekord, Iran
- 3. Graduate Student of Fisheries Science, Faculty of Natural Resources, Isfahan University of Technology, Shahrekord, Iran

*Corresponding author: rasoolzamani@yahoo.com

Received date: 2018.06.10 Reception date: 2019.04.21

Abstract

Algae are among the aquatic organisms that are severely exposed to increasing levels of pollutants, which are becoming a concern in marine ecosystems. Heavy metals are among the compounds for which their increase in high levels causes problems in the structure of algae communities in aquatic ecosystems. In this study, the effect of different concentrations of Nickel (1.5, 3, 4, 6, 8, 12, 15 mg/l) and cobalt (1.5, 2, 3, 5, 6, 8, 10 mg/l) on chlorophyll a and the carotenoid level of microalga Nannochloropsis oculata was investigated for 72 hours on July 2017. The EC₅₀ of nickel and cobalt were determined according to the OECD 2011 protocol. The results of acute tests showed that EC₅₀ of nickel and cobalt was 6.5 mg/l and 5.3 mg/l, respectively. Overall, the results of this study showed that nickel caused toxicity in higher concentrations compared to cobalt. Furthermore, nickel and cobalt heavy metals significantly inhibited the growth and reproduction of N. oculata. However, these heavy metals had lower effect on ceratoid contents.

Keywords: Heavy metals, EC₅₀, Chlorophyll a, Carotenoid, *Nannochloropsis oculata*.