## Assessing the Classification Indices of Ecological Quality Status Using Benthic Macrofauna Identification in Gorgan Bay

Seyed Ghasem Ghorbanzadeh Zafarani'\* Ali Machinchian Moradi' Kazem Darvish Bastami' Seyed MohammadReza Fatemi'

<sup>1</sup>. Research Center for Environment and Sustainable Development (RCESD), Department of Environment, Tehran, Iran.
<sup>r</sup>, <sup>£</sup>. Department of Marine Science, Science and Research Branch, Azad University, Tehran, Iran.
<sup>r</sup>. Iranian National Institute for Oceanography and Atmospheric Science (INIOAS), Tehran, Iran.

\*Corresponding author: ghorbanzadeh

Received date: ۲۰۲۰,۱۱,۱٦ Reception date: ۲۰۲۰,۱۲,۰٦

## Abstract

The present study was conducted for the first time to investigate and identify the ecological quality status (EcoQ) of Gorgan Bay and also to emphasize the degree of agreement between the results of different benthic macrofauna indices, introduction of appropriate index and utilization of basic results in environmental management programs in the region. To do, YY sites were selected and macrobenthic fauna sampling was conducted seasonally in during Y. 17. Totally, 15 species were identified and their biotic indices of BENTIX, BOYA, AMBI and M-AMBI were obtained  $\Upsilon,\Upsilon(\Upsilon-\Upsilon,\circ), \cdot, \varepsilon(\cdot, \cdot, \cdot, \cdot, \cdot, \cdot, \vee)), \Upsilon,\Upsilon(\Upsilon, 1-\Upsilon, \circ),$ and  $\cdot, \mathbf{10}$  ( $\cdot, \mathbf{0}, \cdot, \mathbf{97}$ ), respectively. All the indices indicated a worse EcoQ in eastern part than western part of the bay. A positive and moderate relationship between M-AMBI with Shannon diversity index and BENTIX, as well as between BENTTIX with BOYA exhibited a similar trend in quality assessment of the stations. A partial agreement in EcoQ scores was obtained in  $\gamma\gamma$ ? of the stations while a disagreement was shown in the remaining stations  $(\sqrt[3]{/})$ . However, AMBI and M-AMBI revealed a better and more realistic judgment on EcoQ of the bay which was consistent with the distribution of pollution as an environmental stressor. On the other, since M-AMBI uses the results of AMBI, diversity and richness indices as multimetrics, and more importantly is acquired by the most value of Shannon diversity index as a reference, hence the results of M-AMBI had a stronger judgment on EcoO of the Gorgan bay.

Keywords: M-AMBI, AMBI, BENTIX, BOYA, Gorgan Bay.